

Angiotensin I Converting Enzyme Activity Assay Kit

Cat. No. Kit-2112 **Lot. No.** (See product label)

SPECIFICATION

Product Overview	ACE1 Activity Assay Kit utilizes an active ACE1 to cleave a synthetic o-aminobenzoyl peptide (Abz-based peptide) substrate to release a fluorophore. The released Abz can be easily quantified using a fluorescence microplate reader. This assay kit is simple, rapid and can detect ACE activity as low as 10 mU in biological samples.
Size	100 assays
Description	Angiotensin I converting enzyme (ACE1, EC: 3.4.15.1), a dipeptidyl carboxypeptidase, is part of the renin-angiotensin system (RAS) that controls regulation of blood pressure by cleaving the C-terminal dipeptides of angiotensin I and bradykinin. It is found on the luminal surface of vascular endothelial cells, especially in pulmonary tissues. Elevated levels of ACE1 are found in patients suffering from sarcoidosis, leprosy, hyperthyroidism, acute hepatitis, primary biliary cirrhosis, diabetes mellitus, multiple myeloma, osteoarthritis, amyloidosis, Gaucher's disease, pneumoconiosis, histoplasmosis and miliary tuberculosis.
Applications	Detection of ACE1 activity in tissue/cell lysates Determination of enzymatic activity of purified ACE1
Target Species	Mammalian
Storage	Store kit at -20°C, protected from light. Briefly centrifuge small vials at low speed prior to opening. Read entire protocol before performing the assay. • ACE1 Lysis Buffer and ACE1 Dilution Buffer: Ready to use. Store at -20°C. Bring to room temperature before

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	use. • ACE1 Lysis Buffer: Ready to use. Store at -20°C. Thaw before use. • ACE1 Positive Control: Store at -20°C. Avoid multiple freeze/thaw of the enzyme. Use within 6 months. • ACE1 Substrate: Ready to use. Store at -20°C. Thaw before use. • Abz Standard: Ready to use. Store at -20°C.
Kit Components	ACE1 Assay Buffer: 20 ml ACE1 Lysis Buffer: 40 ml ACE1 Dilution Buffer: 1 ml ACE1 Positive control: 5 µl ACE1 Substrate: 300 µl Abz-Standard (1 mM): 100 µl
Detection method	Fluorometric (Ex/Em = 330/430 nm)
Compatible Sample Types	• Animal tissues: Lung, heart, kidney • Serum, plasma
Features & Benefits	• Simple one-step reaction • Takes only 1-2 hrs • Non-radiometric fluorescent detection • HTP adaptable