

Hexokinase Inhibitor Screening Kit

Cat. No. Kit-2144 **Lot. No.** (See product label)

SPECIFICATION

Product Overview	In the Hexokinase Inhibitor Screening kit, glucose is converted to glucose-6-phosphate by hexokinase; the glucose-6-phosphate is oxidized by glucose-6-phosphate dehydrogenase to form NADH, which reduces a colorless probe to a colored product with strong absorbance at 450 nm. In the presence of a Hexokinase inhibitor, the reaction is impeded/abolished resulting in decrease or total loss of absorbance. This assay kit can be used to screen/study/characterize the potential inhibitors of Hexokinase. The assay is simple, high-throughput adaptable and can be performed within 30 min.
Size	100 assays
Description	Hexokinases (HK) play an important role in glucose metabolism. Hexokinases phosphorylate glucose and generate glucose-6-phosphate for glycolysis. Hexokinases have four isoforms (HK-I, II, III and IV). HK-I, HK-II and HK-III have low Km, while HK-IV has 100 fold high Km. Hexokinase deficiency leads to severe human diseases such as X-linked muscular dystrophy and a rare autosomal recessive hemolytic anemia. On the other hand, increased hexokinase activity is detected in various tumors and is associated with metastasis.
Applications	Screening/characterizing/studying potential inhibitors of Hexokinase
Storage	Store kit at -20°C, protected from light. Briefly centrifuge small vials prior to opening. Read entire protocol before performing the assay. HK Assay Buffer: Bring to room temperature before use. Store at -20°C or 4°C. HK Coenzyme and HK Developer:

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Reconstitute each with 220 μ dH₂O. Pipette up and down to dissolve completely. Store at -20°C. Use within two months. Keep on ice while in use.

HK Converter:
Reconstitute with 220 μ HK Assay Buffer. Aliquot and store at -20°C. Avoid freeze/thaw. Keep on ice while in use. Use within two months.

Hexokinase:
Reconstitute with 160 μ HK Assay Buffer. Aliquot and store at -20°C. Avoid freeze/thaw. Keep on ice while in use. Use within two months.

HK Inhibitor Control:
Reconstitute with 100 μ dH₂O. Aliquot and store at -20°C. Avoid freeze/thaw. Keep on ice while in use. Use within two months.

Kit Components

HK Assay Buffer: 25 ml
HK Substrate: 1 ml
HK Coenzyme: 1 vial
HK Converter: 1 vial
HK Developer: 1 vial
Hexokinase: 1 vial
HK Inhibitor Control: 1 vial

Detection method

Absorbance at 450 nm

Compatible Sample Types

Variable: Inhibitors, small molecules, proteins, etc.

Features & Benefits

- Simple and High throughput adaptable
- Colorimetric Detection
- Screen within 30 min

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA