

Microsome Isolation Kit

Cat. No. Kit-2158 **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Microsome Isolation Kit enables preparation of active microsomes in about one hour, without the need for ultracentrifugation or sucrose gradient fractionation. The kit contains sufficient reagents for 50 isolation procedures, yielding microsomes from roughly 25 grams of tissue or cultured cells.
Size	50 assays
Description	<p>Microsomes are spherical vesicle-like structures formed from membrane fragments following homogenization and fractionation of eukaryotic cells. The microsomal subcellular fraction is prepared by differential centrifugation and consists primarily of membranes derived from the endoplasmic reticulum (ER) and Golgi apparatus. Microsomes isolated from liver tissue are used extensively in pharmaceutical development, toxicology and environmental science to study the metabolism of drugs, organic pollutants and other xenobiotic compounds by the cytochrome P450 monooxidase (CYP) enzyme superfamily. Microsomal preparations are an affordable and convenient in vitro system for assessing Phase I biotransformation reactions, as they contain all of the xenobiotic-metabolizing CYP isozymes and the membrane-bound flavoenzymes (such as NADPH P450-Reductase and cytochrome b5) required for function of the multicomponent P450 enzyme system.</p>
Applications	<p>Convenient and fast isolation of microsomal fraction from animal tissues Assessment of CYP-mediated drug metabolism and xenobiotic biotransformation Protein profiling of microsomal membrane proteins by SDS-PAGE and Western blot</p>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Storage	Store kit at -20°C. Read entire protocol before performing the isolation procedure. Homogenization Buffer and Storage Buffer: Buffers may be stored at -20°C or 4°C. Keep buffers on ice while in use. Protease Inhibitor Cocktail: Resuspend the lyophilized Protease Inhibitor Cocktail in 250 µl anhydrous DMSO (not provided) (500X stock solution) and store at -20°C.
Kit Components	Homogenization Buffer: 80 ml Storage Buffer: 20 ml Protease Inhibitor Cocktail: 1 vial
Compatible Sample Types	<ul style="list-style-type: none">• Mammalian glands and soft tissues such as liver, spleen, lungs etc.• Cultured eukaryotic cell lines such as HepG2 human hepatic carcinoma cells
Features & Benefits	<ul style="list-style-type: none">• Simple & convenient protocol• The kit provides unique formulations of ready-to-use buffers and reagents to isolate microsomes without the need for ultracentrifugation or sucrose gradient fractionation

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA