

Recombinant Human IL19 protein

Cat. No. IL19-130H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Human IL19 protein was expressed in Escherichia coli.
Species	Human
Source	E.coli
ProteinLength	153
Description	Human Interleukin-19 (IL-19) is encoded by the IL19 gene, which is located on the chromosome 1. The protein belongs to the IL-10 family that includes IL-10, IL-20, IL-22, IL-24, and IL-26. As a monomer made of seven amphipathic helices, IL-19 has a helical bundle and shares the same cell surface receptor (IL-20R) with IL-20 and IL-24. It may play some important roles in inflammatory responses because it up-regulates IL-6 and TNF-alpha and induces apoptosis.
Form	Lyophilized from a 0.2µm filtered concentrated solution in PBS, pH 7.4.
Bio-activity	Fully biologically active when compared to standard. The ED50 as determined by a cell proliferation assay using human IL-20Rα and human IL-20Rβ co-transfected murine BaF3 pro-B cells is less than 1.5 ng/ml, corresponding to a specific activity of $> 6.7 \times 10^5$ IU/mg.
Molecular Mass	Approximately 17.9 kDa, a single non-glycosylated polypeptide chain containing 153 amino acids.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

AA Sequence	LRRCLISTDMHHIEESFQEIKRAIQAKDTFPNVTILSTLETLQIIKPLDVCCVTKNLLAFY VDRVFKDHQEPNPKILRKISSIANSFLYMQKTLRQCQEQRQCHCRQEATNATRVIH NYDQLEVHAAAIKSLGELDVFLAWINKNHEVMSSA
Endotoxin	Less than 1 EU/g of rHuIL-19 as determined by LAL method.
Purity	>95% by SDS-PAGE and HPLC analysis.
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in 100 mM HAC to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 centigrade. Further dilutions should be made in appropriate buffered solutions.

GENE INFORMATION

Gene Name	IL19
Official Symbol	IL19
Synonyms	IL19; interleukin 19; interleukin-19; IL 10C; IL 19; MDA1; melanoma differentiation associated protein like protein; NG.1; ZMDA1; melanoma differentiation associated protein-like protein; melanoma differentiation-associated protein-like protein; IL-10C;
Gene ID	29949
mRNA Refseq	NM_013371

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Protein Refseq	NP_037503
MIM	605687
UniProt ID	Q9UHD0

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA