

Active Recombinant Human TNFRSF9 Protein, Fc & Avi-tagged, Biotinylated

Cat. No. TNFRSF9-526H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Biotinylated Recombinant Human TNFRSF9 Protein (Leu 24 - Gln 186) (NP_001552.2), fused with a human IgG1 Fc tag at the C-terminus, followed by a Avi tag, was expressed in HEK293.
Species	Human
Source	HEK293
ProteinLength	Leu 24 - Gln 186
Predicted N Terminal	Leu 24
Form	Lyophilized from sterile 50 mM tris, 100 mM glycine, pH7.5, 10% trehalose.
Bio-activity	Measured by its binding ability in a functional ELISA. Immobilized Human 4?1BB Ligand, Fc Tag at 2µg/mL (100 µL/well) can bind Biotinylated Human 4-1BB, Fc Tag with a linear range of 1-7.8 ng/mL.
Molecular Mass	Recombinant Human TNFRSF9 Protein has a calculated MW of 45.7 kDa. As a result of glycosylation, The protein migrates as 60-66 kDa under reducing (R) condition and 120-130 kDa under non-reducing (NR) condition on SDS-PAGE gel.
Endotoxin	Less than 1.0 EU per µg by the LAL method.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Purity	>95% as determined by SDS-PAGE.
Stability	Samples are stable for up to twelve months from date of receipt at -70°C.
Storage	Store it under sterile conditions at -20°C~-70°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 ug/ul. Centrifuge the vial at 4°C before opening to recover the entire contents.
Conjugation	Biotin

GENE INFORMATION

Gene Name	TNFRSF9 tumor necrosis factor receptor superfamily, member 9 [Homo sapiens]
Official Symbol	TNFRSF9
Synonyms	TNFRSF9; tumor necrosis factor receptor superfamily, member 9; ILA; tumor necrosis factor receptor superfamily member 9; 4 1BB; CD137; CD137 antigen; T cell antigen ILA; T-cell antigen ILA; 4-1BB ligand receptor; homolog of mouse 4-1BB; receptor protein 4-1BB; T-cell antigen 4-1BB homolog; induced by lymphocyte activation (ILA); interleukin-activated receptor, homolog of mouse Ly63; 4-1BB; CDw137; MGC2172; FLJ43501;
Gene ID	3604
mRNA Refseq	NM_001561
Protein Refseq	NP_001552

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

MIM	602250
UniProt ID	Q07011
Chromosome Location	1p36
Pathway	Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem; Downstream signaling in naive CD8+ T cells, organism-specific biosystem;
Function	binding; receptor activity;

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA