

Active Recombinant Human TNFRSF17 Protein, Fc-tagged, FITC-Labeled

Cat. No. TNFRSF17-239H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	FITC-Labeled Recombinant Human TNFRSF17 Protein (Met 1 - Ala 54) (Q02223), fused with a human IgG1 Fc tag at the C-terminus, was expressed in HEK293.
Species	Human
Source	HEK293
ProteinLength	Met 1 - Ala 54
Form	Lyophilized from sterile PBS, pH7.4, 10% trehalose.
Bio-activity	Measured by its binding ability in a functional ELISA. Immobilized Human BAFF, Fc Tag at 10µg/mL (100 µL/well) can bind FITC-Labeled Human BCMA, Fc Tag with a linear range of 0.15-0.625 µg/mL.
Molecular Mass	Recombinant Human TNFRSF17 Protein has a calculated MW of 32 kDa. As a result of glycosylation, The protein migrates as 38-42 kDa under reducing (R) condition and 60-70 kDa under non-reducing (NR) condition on SDS-PAGE gel.
Endotoxin	Less than 1.0 EU per µg by the LAL method.
Purity	>92% as determined by SDS-PAGE.
Stability	Samples are stable for up to twelve months from date of receipt at -70°C.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Storage	Store it under sterile conditions at -20°C~-70°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 ug/ul. Centrifuge the vial at 4°C before opening to recover the entire contents.
Conjugation	FITC

GENE INFORMATION

Gene Name	TNFRSF17 tumor necrosis factor receptor superfamily, member 17 [Homo sapiens]
Official Symbol	TNFRSF17
Synonyms	TNFRSF17; tumor necrosis factor receptor superfamily, member 17; BCMA; tumor necrosis factor receptor superfamily member 17; BCM; CD269; B-cell maturation factor; B cell maturation antigen; B-cell maturation protein;
Gene ID	608
mRNA Refseq	NM_001192
Protein Refseq	NP_001183
MIM	109545
UniProt ID	Q02223
Chromosome Location	16p13.1

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Pathway	Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem; Intestinal immune network for IgA production, organism-specific biosystem; Intestinal immune network for IgA production, conserved biosystem;
Function	receptor activity; tumor necrosis factor-activated receptor activity;

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA