Recombinant Rabbit TIGIT protein, Fc-tagged

Cat. No. TIGIT-260R **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Rabbit TIGIT protein (Ala16-Pro142), fused to human IgG1 Fc tag at C-terminus, was expressed in human 293 cells (HEK293).
Species	Rabbit
Source	HEK293
ProteinLength	127
Description	T-cell immunoreceptor with Ig and ITIM domains (TIGIT) is also known as V-set and immunoglobulin domain-containing protein 9 (VSIG9), V-set and transmembrane domain-containing protein 3 (VSTM3),which belongs to single-pass type I membrane protein containing an immunoglobulin variable domain, a transmembrane domain and an immunoreceptor tyrosine-based inhibitory motif (ITIM). TIGIT is expressed at low levels on peripheral memory and regulatory CD4+ T-cells and NK cells and is up-regulated following activation of these cells (at protein level). TIGIT binds with high affinity to the poliovirus receptor (PVR) which causes increased secretion of IL10 and decreased secretion of IL12B and suppresses T-cell activation by promoting the generation of mature immunoregulatory dendritic cells.
Form	Lyophilized from 0.22 um filtered solution in Tris with Glycine, Arginine and NaCl, pH7.5, 10% trehalose.
Molecular Mass	The protein has a calculated MW of 40.4 kDa. As a result of glycosylation, the protein migrates as 45-50 kDa under reducing (R) condition, and 90-116 kDa under non-



Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing (NR) condition (SDS-PAGE).
Endotoxin	Less than 1.0 EU per ug by the LAL method.
Purity	>95% as determined by SDS-PAGE.
Storage	For long term storage, the product should be stored at lyophilized state at -20 centigrade or lower. Please avoid repeated freeze-thaw cycles. This product is stable after storage at: -20 centigrade to -70 centigrade for 12 months in lyophilized state; -70 centigrade for 3 months under sterile conditions after reconstitution.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 ug/ul. Centrifuge the vial at 4℃ before opening to recover the entire contents.

GENE INFORMATION

Gene Name	TIGIT
Official Symbol	TIGIT
Synonyms	TIGIT; T cell immunoreceptor with Ig and ITIM domains; RGD1563191
Gene ID	100343230
mRNA Refseq	XM_002716684.3
Protein Refseq	XP_002716730.1
UniProt ID	G1SLC1

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

Bioactivity-ELISA of TIGIT-260R	 Immobilized Human Nectin-2, His Tag at 5 μg/mL (100 μL/well) can bind Rabbit TIGIT, Fc Tag with a linear range of 5-78 ng/mL (QC tested).
SDS-PAGE of TIGIT-260R	 Rabbit TIGIT, Fc Tag on SDS-PAGE under reducing (R) and non-reducing (NR) conditions. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95%.

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA