

Active Recombinant Human PDCD1 Protein, His-tagged

Cat. No. PDCD1-190H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human PDCD1 Protein extracellular domain 21-167aa (signal peptide removed), fused with a C-terminal 6*His tag, was expressed in CHO.
Species	Human
Source	CHO
ProteinLength	21-167 a.a.
Description	Programmed cell death 1, also known as PDCD1, is a type I transmembrane glycoprotein, and is an immunoreceptor belonging to the CD28/CTLA-4 family negatively regulates antigen receptor signaling by recruiting protein tyrosine phosphatase, SHP-2 upon interacting with either of two ligands, PD-L1 or PD-L2. PD1 inhibits the T-cell proliferation and production of related cytokines including IL-1, IL-4, IL-10 and IFN-γ by suppressing the activation and transduction of PI3K/AKT pathway. In addition, coligation of PD1 inhibits BCR-mediating signal by dephosphorylating key signal transducer. PD1 has been suggested to be involved in lymphocyte clonal selection and peripheral tolerance, and thus contributes to the prevention of autoimmune diseases. Furthermore, PD1 is shown to be a regulator of virus-specific CD8⁺ T cell survival in HIV infection. As a cell surface molecule, PDCD1 regulates the adaptive immune response. Engagement of PD-1 by its ligands PD-L1 or PD-L2 transduces a signal that inhibits T-cell proliferation, cytokine production, and cytolytic function.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Predicted N Terminal	Pro 21
Form	Lyophilized from sterile PBS, pH 7.4.
Bio-activity	Measured by its binding ability in a functional ELISA. Immobilized human PD-1 at 10 µg/ml (100 µl/well) can bind recombinant human B7-H1 / PD-L1 / Fc chimera with a linear range of 0.02-0.4 µg/ml.
Molecular Mass	The recombinant mature human PD1 consists of 153 amino acids and predicts a molecular mass of 17.4 kDa.
Endotoxin	<1.0EU/ug, LAL method
Purity	>95% (SDS-PAGE)
Stability	Samples are stable for up to twelve months from date of receipt at -70°C.
Storage	Store it under sterile conditions at -20°C~-70°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 ug/ul. Centrifuge the vial at 4°C before opening to recover the entire contents.

GENE INFORMATION

Gene Name	PDCD1 programmed cell death 1 [Homo sapiens]
Official Symbol	PDCD1
Synonyms	PDCD1; programmed cell death 1; programmed cell death protein 1; CD279; PD1; protein PD-1; PD-1; SLEB2; hPD-1; hPD-I;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene ID	5133
mRNA Refseq	NM_005018
Protein Refseq	NP_005009
MIM	600244
UniProt ID	Q15116
Chromosome Location	2q37.3
Pathway	Adaptive Immune System, organism-specific biosystem; Cell adhesion molecules (CAMs), organism-specific biosystem; Cell adhesion molecules (CAMs), conserved biosystem; Costimulation by the CD28 family, organism-specific biosystem; Immune System, organism-specific biosystem; PD-1 signaling, organism-specific biosystem; T cell receptor signaling pathway, organism-specific biosystem;
Function	protein binding; protein tyrosine phosphatase activity; signal transducer activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA