

Active Recombinant Human TNFSF9 Protein, Fc-tagged

Cat. No. TNFSF9-719H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human TNFSF9 Protein extracellular domain 51-254aa (signal peptide removed), fused with a N-terminal Fc tag, was expressed in HEK293.
Species	Human
Source	HEK293
ProteinLength	51-254 a.a.
Description	4-1BBL is the high affinity ligand of 4-1BB, also known as CD137L or TNFSF9. It is shown to be a type II surface glycoprotein belonging to the TNF superfamily. Expression of 4-1BBL is restricted to APCs, such as dendritic cells, macrophages, and activated B cells. Members of the TNFTNF receptor superfamily have been shown to play critical roles in regulating cellular activation, differentiation and apoptosis. Several studies have reported that 4-1BBL/4-1BB interaction provided a co-stimulatory signal to T cells, and increased T cell proliferation and cytokines production. In addition, 4-1BBL is involved in cancers, infectious diseases and autoimmune diseases.
Form	Lyophilized from sterile PBS, pH 7.4.
Bio-activity	Measured by its binding ability in a functional ELISA.
Molecular Mass	The recombinant mature human CD137L protein consists of 436 amino acids and has a calculated molecular mass of 47.4 KD.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Endotoxin	<1.0EU/ug, LAL method
Purity	>95% (SDS-PAGE)
Stability	Samples are stable for up to twelve months from date of receipt at -70°C.
Storage	Store it under sterile conditions at -20°C~-70°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 ug/ul. Centrifuge the vial at 4°C before opening to recover the entire contents.

GENE INFORMATION

Gene Name	TNFSF9 tumor necrosis factor (ligand) superfamily, member 9 [Homo sapiens]
Official Symbol	TNFSF9
Synonyms	TNFSF9; tumor necrosis factor (ligand) superfamily, member 9; tumor necrosis factor ligand superfamily member 9; 4 1BB L; homolog of mouse 4 1BB L; receptor 4 1BB ligand; 4-1BBL; 4-1BB ligand; receptor 4-1BB ligand; homolog of mouse 4-1BB-L; CD137L; 4-1BB-L;
Gene ID	8744
mRNA Refseq	NM_003811
Protein Refseq	NP_003802
MIM	606182

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID	P41273
Chromosome Location	19p13.3
Pathway	Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem;
Function	cytokine activity; receptor binding; tumor necrosis factor receptor binding;

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA