

Recombinant HIV HIVTPO41 protein, β-gal-tagged

Cat. No. HIVTPO41-161H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	HIV Type-O gp41 recombinant, containing 250 a.a. of the HIV Type-O immunodominant regions from gp41 protein is fused with a Beta-galactosidase at N-terminus having a total Mw of 94kDa.
Species	HIV
Source	E.coli
ProteinLength	250 a.a.
Description	Human immunodeficiency virus (HIV) is a retrovirus that can lead to a condition in which the immune system begins to fail, leading to opportunistic infections. HIV primarily infects vital cells in the human immune system such as helper T cells (specifically CD4+ T cells), macrophages and dendritic cells. HIV infection leads to low levels of CD4+ T cells through three main mechanisms: firstly, direct viral killing of infected cells; secondly, increased rates of apoptosis in infected cells; and thirdly, killing of infected CD4+ T cells by CD8 cytotoxic lymphocytes that recognize infected cells. When CD4+ T cell numbers decline below a critical level, cell-mediated immunity is lost, and the body becomes progressively more susceptible to opportunistic infections. HIV was classified as a member of the genus Lentivirus, part of the family of Retroviridae. Lentiviruses have many common morphologies and biological properties. Many species are infected by lentiviruses, which are characteristically responsible for long-duration illnesses with a long incubation period. Lentiviruses are transmitted as single-stranded, positive-sense, enveloped RNA

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

viruses. Upon entry of the target cell, the viral RNA genome is converted to double-stranded DNA by a virally encoded reverse transcriptase that is present in the virus particle. This viral DNA is then integrated into the cellular DNA by a virally encoded integrase so that the genome can be transcribed. Once the virus has infected the cell, two pathways are possible: either the virus becomes latent and the infected cell continues to function, or the virus becomes active and replicates, and a large number of virus particles are liberated that can then infect other cells.

Form	Sterile filtered colorless clear solution. 1.5M urea, 25mM Tris-HCl pH 8.0 and 50% Glycerol.
Molecular Mass	94kDa
Purity	Greater than 90.0% as determined by SDS-PAGE.
Usage	The products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.
Stability	HIV Type-O although stable at 4°C for 1 week, should be stored below -18°C. Please prevent freeze thaw cycles.
Specificity	Immunoreactive with all sera of HIV type-O infected individuals.
Shipping	Ice Packs