

Recombinant HIV env protein, β-gal-tagged

Cat. No. env-168H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	The E.Coli derived protein contains the full-length sequence of N-terminal epitopes of HIV-I gp41 395 amino acids (444-833) immunodominant regions gp41L. The protein is fused to b-galactosidase (114kDa) at N-Terminus.
Species	HIV
Source	E.coli
Description	Human immunodeficiency virus (HIV) is a retrovirus that can lead to a condition in which the immune system begins to fail, leading to opportunistic infections. HIV primarily infects vital cells in the human immune system such as helper T cells (specifically CD4+ T cells), macrophages and dendritic cells. HIV infection leads to low levels of CD4+ T cells through three main mechanisms: firstly, direct viral killing of infected cells; secondly, increased rates of apoptosis in infected cells; and thirdly, killing of infected CD4+ T cells by CD8 cytotoxic lymphocytes that recognize infected cells. When CD4+ T cell numbers decline below a critical level, cell-mediated immunity is lost, and the body becomes progressively more susceptible to opportunistic infections. HIV was classified as a member of the genus Lentivirus, part of the family of Retroviridae. Lentiviruses have many common morphologies and biological properties. Many species are infected by lentiviruses, which are characteristically responsible for long-duration illnesses with a long incubation period. Lentiviruses are transmitted as single-stranded, positive-sense, enveloped RNA viruses. Upon entry of the target cell, the viral RNA genome is converted to double-stranded DNA by a virally encoded reverse transcriptase that is present in the virus.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

particle. This viral DNA is then integrated into the cellular DNA by a virally encoded integrase so that the genome can be transcribed. Once the virus has infected the cell, two pathways are possible: either the virus becomes latent and the infected cell continues to function, or the virus becomes active and replicates, and a large number of virus particles are liberated that can then infect other cells.

Form	Sterile filtered colorless clear solution. 8M Urea, 20mM Tris-HCl pH 8.0, 10mM b-mercaptoethanol.
AA sequence	IEFPGIFRPGGGDMRDNRSELYKYKVVKIEPLGVAPTKAKRRVVQREKRAVGIGAL FLGFLGAAGSTMGAASMTLTVQARQLLSGIVQQQNNLLRAIEAQQHLLQLTVWGIKQ LQARILAVERYLKDQQLLGIWGC SGKLICTTAVPWNASWSNKSLEQIWNMTWME WDREINNYTSLIHSLIEESQNQQEKNEQELLELDKWASLWNWFNITNWLWYIKLFIMI VGGLVGLRIVFAVLSVVRQGYSPFSFQTHLPIPRGPDRPEGIEEEGGERDRDRS IRLVNGSLALIWDLRLSLCLFSYHRLRDLLLIVTRIVELLGRRGWEALKYWWNLLQYW SQELKNSAVSLLNATAIAVAEGTDRVIEVVQGAYRAIRHIPRRIRQGLERILL
Purity	Greater than 95.0% as determined by HPLC analysis & SDS-PAGE.
Usage	The products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.
Stability	HIV-1 gp41 Long although stable at 4°C for 1 week, should be stored below -18°C. Please prevent freeze thaw cycles.
Specificity	Immunoreactive with all sera of HIV-1 infected individuals.
Shipping	Ice Packs

GENE INFORMATION

Gene Name	env gp160; envelope glycoprotein [Human immunodeficiency virus 1]
Official Symbol	env
Synonyms	env; gp160; envelope glycoprotein; Envelope surface glycoprotein gp160, precursor; hypothetical protein; Envelope surface glycoprotein gp120; Envelope transmembrane domain
Gene ID	155971
Protein Refseq	NP_057856.1
UniProt ID	P04578

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA