

Recombinant HIV HIVP14 protein, β-gal-tagged

Cat. No. HIVP14-181H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	The E.coli derived 39kDa recombinant protein is a non-glycosylated polypeptide chain, containing the HIV-1 p24 gag immunodominant regions, 77-436 amino acids. The HIV-1 p24 gag is fused to beta-galactosidase (114kDa).
Species	HIV
Source	E.coli
ProteinLength	77-436 amino acids
Description	Human immunodeficiency virus (HIV) is a retrovirus that can lead to a condition in which the immune system begins to fail, leading to opportunistic infections. HIV primarily infects vital cells in the human immune system such as helper T cells (specifically CD4+ T cells), macrophages and dendritic cells. HIV infection leads to low levels of CD4+ T cells through three main mechanisms: firstly, direct viral killing of infected cells; secondly, increased rates of apoptosis in infected cells; and thirdly, killing of infected CD4+ T cells by CD8 cytotoxic lymphocytes that recognize infected cells. When CD4+ T cell numbers decline below a critical level, cell-mediated immunity is lost, and the body becomes progressively more susceptible to opportunistic infections. HIV was classified as a member of the genus Lentivirus, part of the family of Retroviridae. Lentiviruses have many common morphologies and biological properties. Many species are infected by lentiviruses, which are characteristically responsible for long-duration illnesses with a long incubation period. Lentiviruses are transmitted as single-stranded, positive-sense, enveloped RNA

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

viruses. Upon entry of the target cell, the viral RNA genome is converted to double-stranded DNA by a virally encoded reverse transcriptase that is present in the virus particle. This viral DNA is then integrated into the cellular DNA by a virally encoded integrase so that the genome can be transcribed. Once the virus has infected the cell, two pathways are possible: either the virus becomes latent and the infected cell continues to function, or the virus becomes active and replicates, and a large number of virus particles are liberated that can then infect other cells.

Form

Sterile filtered colorless clear solution. 8M urea, 20mM Tris-HCl pH-8 & 10mM β-mercaptoethanol.

Molecular Mass

39kDa

AA sequence

slyntvatlycvhqrieikdtkealdkikeeqnkskkkaqqaadtghssqvsqnyqivqniqqgmvhqaisprtlhaw
vkvveekafspevipmfsalsegatpqdlntmlntvgghqaamqmlketineeaaewdrvhpvhagpiapgqmr
eprgsdiagtstlqeqigwmtnnppipvgeiykrwiilgnkivrmysptsildirgpkpfrdyvdrfykltlaeqasqe
vknwmtetllvqnanpdcktilkalgpaatleemmtacqvggpgghkarvlaeamsqvtnsatimmqrgnfrnqrki
vkcfncgkeghiarncraprkkgcwkcqgkeghqmkdcterqanflgk.

Purity

Greater than 95.0% as determined by HPLC analysis and SDS-PAGE.

Applications

HIV-1 p24 gag antigen in ELISA and Western blots, excellent antigen for early detection of HIV seroconvertors with minimal specificity problems.

Usage

The products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Stability

HIV-1 p24 gag although stable at 4°C for 1 week, should be stored below -18°C. Please prevent freeze thaw cycles.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Specificity	Immunoreactive with all sera of HIV-1 infected individuals.
--------------------	---

Shipping	Ice Packs
-----------------	-----------

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA