

GMP Recombinant Human PDGFB protein

Cat. No. PDGFB-4346HG **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human PDGFB protein was expressed in Escherichia coli. This product is produced under cGMP guidelines.
Species	Human
Source	E.coli
ProteinLength	110
Description	Platelet-derived growth factor (PDGF) presenting in serum but absent from plasma was first discovered in animal study by Lynch and co-workers in the late 1980s. It is a disulfide-linked dimer consisting of two peptides-chain A and chain B. PDGF has three subforms: PDGF-AA, PDGF-BB, PDGF-AB. It is involved in a number of biological processes, including hyperplasia, embryonic neuron development, chemotaxis, and respiratory tubule epithelial cell development. The function of PDGF is mediated by two receptors (PDGFR- α and PDGFR- β).
Form	Lyophilized from a 0.2 μ m filtered concentrated solution in PBS, pH 7.4.
Bio-activity	Fully biologically active when compared to standard. The ED50 as determined by a cell proliferation assay using murine Balb/c 3T3 cells is less than 3 ng/ml, corresponding to a specific activity of $> 3.3 \times 10^5$ IU/mg.
Molecular Mass	Approximately 24.8 kDa, a disulfide-linked homodimeric protein containing two 110 amino acid residues polypeptide (B chain).

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

AA Sequence	MSLGSLTIAEPAMIAECKTRTEVF EISRRLIDRTNANFLVWPPCVEVQRCSGCCNNR NVQCRPTQVQLRPVQVRKIEIVRKKPIFKKATVTLEDHLACKCETVAAARPVT
Endotoxin	Less than 0.01 EU/μg of rHuPDGF-BB GMP as determined by LAL method.
Purity	>98% by SDS-PAGE and HPLC analysis.
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤-20 centigrade. Further dilutions should be made in appropriate buffered solutions.

GENE INFORMATION

Gene Name	PDGFB
Official Symbol	PDGFB
Synonyms	PDGFB; platelet-derived growth factor beta polypeptide; platelet derived growth factor beta polypeptide (simian sarcoma viral (v sis) oncogene homolog) , SIS; platelet-derived growth factor subunit B; becaplermin; oncogene SIS; SSV; PDGF-2; PDGF, B chain; PDGF subunit B; proto-oncogene c-Sis; platelet-derived growth factor 2; platelet-derived growth factor B chain; platelet-derived growth factor, B chain; Platelet-derived growth factor, beta polypeptide (oncogene SIS); platelet-derived growth factor beta polypeptide (simian sarcoma viral (v-sis) oncogene homolog); SIS;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

PDGF2; c-sis; FLJ12858;

Gene ID 5155


mRNA Refseq NM_002608

Protein Refseq NP_002599

MIM 190040

UniProt ID P01127

**SDS-PAGE of
PDGFB-4346HG**




 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

HPLC of PDGFB- 4346HG



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA