

Active Recombinant Human IL17RC protein, Fc-tagged

Cat. No. IL17RC-249H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human IL17RC(Leu21-His465) fused Fc region of Human IgG1 at C-terminal was expressed in HEK293.
Species	Human
Source	HEK293
ProteinLength	21-465 a.a.
Description	IL-17 receptor C (IL-17RC; also known as IL-17RL and IL-17Rhom) is an 85-110 kDa member of the IL-17 receptor family. IL-17 plays a key role in the development of autoimmunity and inflammation. Human IL-17 RC is a type I transmembrane glycoprotein that is expressed on neutrophils, endothelial cells, chondrocytes and osteoblasts, breast and prostatic epithelium, fibroblasts, renal tubular epithelium, and skeletal muscle cells. Mature human IL-17 RC consists of a 518 amino acid (aa) extracellular domain (ECD), a 21 aa transmembrane segment, and a 232 aa cytoplasmic region. It is estimated that there are over 90 alternative splice forms, with transmembrane-containing isoforms predominating. This recombinant protein corresponds to IL-17 RC isoform 1, the canonical form. Within the ECD, IL-17 RC isoform 1 shares 69% aa sequence identity with mouse IL-17 RC. IL-17 RC is the cognate receptor for IL-17F, and it binds IL-17A with similar affinity. IL-17 RC associates with IL-17 RA to form a receptor complex for both IL-17A and IL-17F.
Predicted N Terminal	Leu21

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Form	Lyophilized from a 0.2 µm filtered solution in PBS.
Bio-activity	Measured by its binding ability in a functional ELISA. When Recombinant Human IL-17RC Isoform 1 Fc Chimera is coated at 0.5 µg/mL, Recombinant Human IL-17/IL-17A binds with an ED50 = 0.1-0.6 ng/mL.
Molecular Mass	Predicted Molecular Mass: 76 kDa SDS-PAGE: 95-113 kDa, reducing conditions
Endotoxin	<0.1 EU per 1 µg of the protein by the LAL method.
Purity	>95%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.
Reconstitution	Reconstitute at 500 µg/mL in PBS.
GENE INFORMATION	
Gene Name	IL17RC interleukin 17 receptor C [Homo sapiens]
Official Symbol	IL17RC
Synonyms	IL17RC; interleukin 17 receptor C; interleukin-17 receptor C; IL17 RL; IL-17RL; IL17Rhom; zcytoR14; IL17F receptor; IL-17 receptor C; interleukin-17 receptor homolog; interleukin-17 receptor-like protein; IL17RL; IL17-RL; FLJ95963; FLJ96005; MGC10763;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene ID	84818
mRNA Refseq	NM_001203263
Protein Refseq	NP_001190192
MIM	610925
UniProt ID	Q8NAC3

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA