

Active Recombinant Mouse Pdgfa protein, His-tagged

Cat. No. Pdgfa-564M **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Active Recombinant Mouse Pdgfa protein(P20033)(90~190 aa), fused with N-terminal His tag, was expressed in E.coli.
Species	Mouse
Source	E.coli
ProteinLength	90-190 aa
Form	Lyophilized from sterile 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Bio-activity	PDGFA (Platelet-derived growth factor subunit A) is a Growth factor that plays an essential role in the regulation of embryonic development, cell proliferation, cell migration, survival and chemotaxis. PDGFA has been described as a chemoattractant for monocytes and proven to be able to induce chemotactic migration of THP-1 cells. Therefore, chemotaxis assay used 24-well microchemotaxis system was undertaken to detect the chemotactic effect of PDGFA on the human monocytic cell line THP-1. Briefly, THP-1 cells were seeded into the upper chambers (100uL cell suspension, 106cells/mL in RPMI 1640 with 0.5% FBS) and PDGFA (1.5ng/mL, 7.5ng/mL and 15ng/mL diluted separately in serum free RPMI 1640) was added in lower chamber with a polycarbonate filter (8µm pore size) used to separate the two compartments. After incubation at 37°C with 5%CO2 for 3h, the filter was removed, then cells in low chamber were observed by inverted microscope at low magnification

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

(×40) and the number of migrated cells were counted at high magnification (×400) randomly (five fields for each filter). Result shows PDGFA is able to induce migration of THP-1 cells. The migrated THP-1 cells in low chamber at low magnification (×40) were shown. Five fields of each chamber were randomly chosen, and the migrated cells were counted at high magnification (×400). Statistical results were shown. The optimum chemotaxis of PDGFA occurs at 1.5ng/mL.

Molecular Mass	15.2kDa
Endotoxin	<1.0EU per 1µg (determined by the LAL method).
Purity	> 97% as determined by SDS-PAGE.
Storage	Avoid repeated freeze/thaw cycles. Store at -20°C for 12 months. Aliquot and store at -80°C for 12 months.
Reconstitution	Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

GENE INFORMATION

Gene Name	Pdgfa platelet derived growth factor, alpha [Mus musculus]
Official Symbol	Pdgfa
Synonyms	PDGFA; platelet derived growth factor, alpha; platelet-derived growth factor subunit A; PDGF-1; PDGF subunit A; platelet-derived growth factor A chain; platelet-derived growth factor alpha polypeptide;
Gene ID	18590

 Tel: 1-631-559-9269 1-516-512-3133


 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA


mRNA Refseq NM_008808

Protein Refseq NP_032834

Activity



SDS-PAGE



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA