

Native Canavalia ensiformis Lectin, FITC conjugated

Cat. No. Lectin-1723C **Lot. No.** (See product label)

SPECIFICATION

Product Overview	This product is canavalia ensiformis lectin which against alpha-linked mannose and glucose (FITC). Affinity purified ConA is conjugated with Fluorescein-5-isothiocyanate (FITC-isomer 1) and purified on Gel filtration column, A max. = 490 nm, E max =520 nm, Ratio of A490/A280= 1.0-1.4.
Species	Canavalia ensiformis
Source	Canavalia ensiformis
Description	Concanavalin A is a lectin protein (MW 104kDa), homotetramer 26 kDa; originally extracted from the jack-bean, Canavalia ensiformis. It binds specifically to certain structures found in various sugars α-mannosyl and α-glucosyl residues in glycoproteins. It was the first lectin to be available on a commercial basis and is widely used in biology and biochemistry to characterize glycoproteins and other sugar-containing entities. It is also used to purify macromolecules in lectin affinity chromatography. Concanavalin A interacts with diverse receptors containing mannose carbohydrates (serum and membrane glycoproteins).ConA agglutinate strongly erythrocytes without being blood group specific. Normal cell re agglutinated after trypsinisation. ConA is a also a lymphocyte mitogen.ConA reacts with many bacteria, like E. coli Dictyostelium discoideum et B. subtilis It is also widely believed to be involved in the interaction between alpha-mannosyl oligosaccharides on the surface of the HIV virus and the human T cell lymphocyte. Inhibiting/Eluting sugars: 200 α-mM α-methyl mannoside / 200 mM α-methyl glucoside mixture.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	Carbohydrate-Binding Specificity of Con A: (Man α 1,2Man α 1,2Man > Man α 1,2Man > α -Man > α -Glc > α GlcNAc)
Form	10 mM bicarbonate, 150 mM NaCl, pH 8, 0.1 mM Calcium chloride, 0.01mM manganese chloride and 0.05% sodium azide.
Applications	ICC/IF. Immunofluorescence 1:250-1:1000. The optimum dilution should be determined by the individual lab. Con A has an Isoelectric point of about pH 4.5-5.5 and requires calcium or manganese ions at each of its four saccharide binding sites, these ions should be used in buffer. Do not use phosphate buffer for dilution of this lectin as it will decrease the activity of lectin.
Storage	Store at 2-8centigrade.
Concentration	5 mg/ml (Please refer to the vial label for the specific concentration)
GENE INFORMATION	
Synonyms	Concanavalin A, Concanavalin A, Canavalia ensiformis, Canavalia ensiformis lectin, ConA lectin, Canavalia ensiformis, ConA, Concanavalin A lectin, Con A lectin
UniProt ID	P02866