

Recombinant Human XRCC6 protein

Cat. No. XRCC6-1548H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human XRCC6 protein was expressed in E. coli.
Species	Human
Source	E.coli
Description	Ku70 forms a protein complex with Ku80. The Ku70-Ku80 complex binds to DNA ends and doublestranded DNA breaks. This complex is involved in nonhomologous end joining (NHEJ) for double-strand DNA break repair, and is also involved in NHEJ during V(D)J recombination of immunoglobulin genes.
Form	50 mM Tris-HCl (pH 7.5), 5 mM β-mercaptoethanol, 500 mM NaCl, and 10% glycerol.
Molecular Mass	70 kDa

GENE INFORMATION

Gene Name	XRCC6 X-ray repair complementing defective repair in Chinese hamster cells 6 [Homo sapiens]
Official Symbol	XRCC6
Synonyms	XRCC6; X-ray repair complementing defective repair in Chinese hamster cells 6; G22P1, thyroid autoantigen 70kD (Ku antigen) , thyroid autoantigen 70kDa (Ku antigen); X-ray repair cross-complementing protein 6; D22S671; D22S731; Ku

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

autoantigen; 70kDa; KU70; ML8; 5-dRP lyase Ku70; Ku autoantigen, 70kDa; DNA repair protein XRCC6; Ku autoantigen p70 subunit; 70 kDa subunit of Ku antigen; thyroid-lupus autoantigen p70; lupus Ku autoantigen protein p70; 5-deoxyribose-5-phosphate lyase Ku70; CTC box binding factor 75 kDa subunit; CTC box-binding factor 75 kDa subunit; thyroid autoantigen 70kD (Ku antigen); ATP-dependent DNA helicase 2 subunit 1; thyroid autoantigen 70kDa (Ku antigen); ATP-dependent DNA helicase II 70 kDa subunit; ATP-dependent DNA helicase II, 70 kDa subunit; TLAA; CTC75; CTCBF; G22P1;

Gene ID	2547
mRNA Refseq	NM_001469
Protein Refseq	NP_001460
MIM	152690
UniProt ID	P12956