

Active Recombinant Human ANGPT2 protein, His-tagged

Cat. No. ANGPT2-531H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Active Recombinant Human ANGPT2 protein(O15123)(24~165 aa), fused with N-terminal His tag, was expressed in E.coli.
Species	Human
Source	E.coli
ProteinLength	24-165 aa
Form	Lyophilized from sterile 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Bio-activity	Angiopoietin is part of a family of vascular growth factors that play a role in embryonic and postnatal angiogenesis. Angiopoietin cytokines are involved with controlling microvascular permeability, vasodilation, and vasoconstriction by signaling smooth muscle cells surrounding vessels. There are now four identified angiopoietins: ANGPT1, ANGPT2, ANGPT3, ANGPT4. Angiopoietin 2 (ANGPT2) promotes cell death and disrupts vascularization. Yet, when it is in conjunction with vascular endothelial growth factors, or VEGF, it can promote neo-vascularization. Besides, TEK Tyrosine Kinase (Tie2) has been identified as an interactor of ANGPT2, thus a binding ELISA assay was conducted to detect the interaction of recombinant human ANGPT2 and recombinant humanTie2. Briefly, ANGPT2 were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100uL were then transferred to Tie2-coated microtiter wells and incubated for 2h at 37°C. Wells were washed with PBST

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

and incubated for 1h with anti-ANGPT2 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50μ stop solution to the wells and read at 450nm immediately. The binding activity of ANGPT2 and Tie2 was in a dose dependent manner.

Molecular Mass	17.6kDa
Purity	> 95% as determined by SDS-PAGE.
Storage	Avoid repeated freeze/thaw cycles. Store at -20°C for 12 months. Aliquot and store at -80°C for 12 months.
Reconstitution	Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

GENE INFORMATION

Gene Name	ANGPT2 angiopoietin 2 [Homo sapiens]
Official Symbol	ANGPT2
Synonyms	ANGPT2; angiopoietin 2; angiopoietin-2; Ang2; ANG-2; Tie2-ligand; angiopoietin-2B; angiopoietin-2a; ANG2; AGPT2;
Gene ID	285
mRNA Refseq	NM_001118887
Protein Refseq	NP_001112359

 Tel: 1-631-559-9269 1-516-512-3133


 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA


MIM 601922

UniProt ID O15123

Activity



SDS-PAGE



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA