

Active Recombinant Human CX3CL1 protein, His-tagged

Cat. No. CX3CL1-765H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Active Recombinant Human CX3CL1 protein(P78423)(81~336 aa), fused with N-terminal His tag, was expressed in E.coli.
Species	Human
Source	E.coli
ProteinLength	81-336 aa
Form	Lyophilized from sterile PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
Bio-activity	Chemokine C-X3-C-Motif Ligand 1 (CX3CL1) also known as fractalkine is a large cytokine protein of 373 amino acids, it contains multiple domains and is the only known member of the CX3C chemokine family. Soluble CX3CL1 potently chemoattracts T cells and monocytes, while the cell-bound chemokine promotes strong adhesion of leukocytes to activated endothelial cells, where it is primarily expressed. Thus, chemotaxis assay used 24-well microchemotaxis system was undertaken to detect the chemotactic effect of CX3CL1 on the human monocytic cell line THP-1. Briefly, THP-1 cells were seeded into the upper chambers (150ul cell suspension, 10^6 cells/ml in RPMI 1640 with FBS free) and CX3CL1 (0.1ng/ml, 0.01ng/ml, 0.001ng/ml, 0.0001 and 0.00001ng/ml diluted separately in serum free RPMI 1640) was added in lower chamber with a polycarbonate filter(8um pore size) used to separate the two compartments. After incubation at 37°C with 5% CO ₂ for 1h,

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

the filter was removed, then cells in low chamber were observed by inverted microscope at low magnification ($\times 10$) and the number of migrated cells were counted using Fluorescence Activating Cell Sorter. Result shows CX3CL1 is able to induce migration of THP-1 cells. The migrated THP-1 cells in low chamber at low magnification ($\times 10$) were shown. Statistical results of FACS were shown. The optimum chemotaxis of CX3CL1 occurs at 0.001-0.00001 ng/ml.

Molecular Mass	28.5kDa
Endotoxin	<1.0EU per 1 μ g (determined by the LAL method).
Purity	> 95% as determined by SDS-PAGE.
Storage	Avoid repeated freeze/thaw cycles. Store at -20°C for 12 months. Aliquot and store at -80°C for 12 months.
Reconstitution	Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.


GENE INFORMATION

Gene Name	CX3CL1 chemokine (C-X3-C motif) ligand 1 [Homo sapiens]
Official Symbol	CX3CL1
Synonyms	CX3CL1; chemokine (C-X3-C motif) ligand 1; SCYD1, small inducible cytokine subfamily D (Cys X3 Cys), member 1 (fractalkine, neurotactin); fractalkine; ABCD 3; C3Xkine; CXC3; CXC3C; neurotactin; NTN; C-X3-C motif chemokine 1; small-inducible cytokine D1; CX3C membrane-anchored chemokine; small inducible cytokine subfamily D (Cys-X3-Cys), member-1; small inducible cytokine subfamily D (Cys-X3-Cys), member 1 (fractalkine, neurotactin); NTT; SCYD1; ABCD-3;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene ID	6376
mRNA Refseq	NM_002996
Protein Refseq	NP_002987
MIM	601880
UniProt ID	P78423
Activity	
SDS-PAGE	