

Active Native Ricinus Communis Agglutinin I Protein, Rhodamine labeled

Cat. No. Lectin-1731R **Lot. No.** (See product label)

SPECIFICATION

Product Overview	This product is the Ricinus Communis Agglutinin I and has sugar specificity against galactose and lactose. The excitation maximum is at 550 nm and the emission maximum is at 575 nm.
Species	Ricinus Communis
Source	Ricinus Communis
Description	This lectin consists of two subunits of 60 kDa which can be dissociated by reducing agents into closely related chains between 27 kDa and 33 kDa. One of the chains appears to be common to the "B" chain of another castor bean lectin, ricin, while the other chain is unique to RCA I. The B chain binds to galactose or N-acetylgalactosamine residues of membrane glycoconjugates.
Form	10 mM HEPES, 0.15 M NaCl, pH 7.5, 0.08% sodium azide, 0.1 mM Ca ²⁺
Bio-activity	Carbohydrate-Binding Specificity: Galactose, Lactose.
Molecular Mass	120 kDa
Applications	Immunofluorescence, Glycobiology
Usage	The recommended concentration range for use is 5-20 μ g/ml. It has been found to have moderate toxicity in some animal models, but its toxicity to humans has not

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

been fully determined, and a related protein, RCA 60, is highly toxic to humans. Never pipet solutions of this lectin by mouth. DO NOT LYOPHILIZE!

Storage

Refrigerate in the dark. A precipitate may form during storage. This does not have a significant adverse effect on the product. If a precipitate forms upon long-term storage at 4 centigrade, warm to 37 centigrade. and centrifuge before use.

Concentration

5 mg/ml

GENE INFORMATION**Synonyms**

Lectin; RCA I; RCA120

UniProt ID

[P02879](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA