

Active Native *Canavalia ensiformis* Concanavalin A Protein, Rhodamine labeled

Cat. No. Lectin-1736C **Lot. No.** (See product label)

SPECIFICATION

Product Overview	This product is the <i>Canavalia ensiformis</i> (Jack Bean) seeds Concanavalin A and has sugar specificity against Mannose and Glucose. The excitation maximum is at 550 nm and the emission maximum is at 575 nm.
Species	<i>Canavalia ensiformis</i> (Jack Bean) seeds
Source	<i>Canavalia ensiformis</i> (Jack Bean) seeds
Description	Con A recognizes α -linked mannose present as part of a core oligosaccharide in many serum and membrane glycoproteins. At neutral and alkaline pH, Con A exists as a tetramer of four identical subunits; below pH 5.6, Con A dissociates into active dimers of 52 kDa. Acetylation, succinylation, or other derivatizations can also produce stable forms with dimeric structures. (See succinylated Con A). Nicks in the sequence are often present in the purest preparations due to hydrolytic damage within the seeds.
Form	10 mM HEPES, 0.15 M NaCl, pH 7.5, 0.1 mM Ca^{2+} 0.08% sodium azide, 0.01 mM Mn^{2+}
Bio-activity	Inhibiting/Eluting Sugar: mixture of 200 mM α -methylmannoside/200 mM α -methylglucoside
Molecular Mass	104 kDa

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications	Immunofluorescence, Glycobiology
Usage	The recommended concentration range for use is 5-20 μ g/ml.
Storage	Refrigerate in the dark. If precipitate forms upon long term storage at 4 centigrade, warm to 37 centigrade. and centrifuge before use. This does not have a significant adverse effect on the product.
Concentration	5.0 mg/ml active conjugate
GENE INFORMATION	
Synonyms	Lectin; Con A
UniProt ID	P02866

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA