

Active Native Wheat Germ Agglutinin Protein, Peroxidase conjugated

Cat. No. Lectin-1737W **Lot. No.** (See product label)

SPECIFICATION

Product Overview	This product is the Wheat germ Agglutinin and has sugar specificity against N-Acetylglucosamine. The peroxidase-wheat germ agglutinin is specially prepared for neuronal transport studies.
Species	Wheat germ
Source	Wheat Germ
Description	Wheat germ agglutinin (WGA) contains a group of closely related isolectins, with an isoelectric point about pH 9. The receptor sugar for WGA is N-acetylglucosamine, with preferential binding to dimers and trimers of this sugar. WGA can bind oligosaccharides containing terminal N-acetylglucosamine or chitobiose, structures which are common to many serum and membrane glycoproteins. Bacterial cell wall peptidoglycans, chitin, cartilage glycosaminoglycans, and glycolipids can also bind WGA. Native WGA has also been reported to interact with some glycoproteins via sialic acid residues (see succinylated WGA). This lectin is used for the purification of insulin receptors and for neuronal tracing.
Form	Buffered salt, 0.02% Thimerosal
Bio-activity	Inhibiting/Eluting Sugar: Chitin Hydrolysate or 500 mM N-acetylglucosamine with salt and/or acid elution generally required.
Molecular Mass	36 kDa

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications	Immunohistochemistry; Immunocytochemistry; Neurobiology; Blotting Applications; ELISAs; Glycobiology
---------------------	--

Storage	Refrigerate in the dark.
----------------	--------------------------

Concentration	40 mg/ml
----------------------	----------

GENE INFORMATION

Synonyms	Lectin; WGA
-----------------	-------------

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA