

Recombinant Human PPIG, His-tagged

Cat. No. PPIG-28674TH **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant fragment, corresponding to amino acids 1-175 of Human PPIG fused to a His tag at the N terminus; 195 aa, 21.6 kDa.
Species	Human
Source	E.coli
ProteinLength	1-175 a.a.
Description	Peptidyl-prolyl cis-trans isomerase G is an enzyme that in humans is encoded by the PPIG gene.
Conjugation	HIS
Tissue specificity	Ubiquitous.
Form	Liquid
Purity	>95% by SDS-PAGE
Storage buffer	Preservative: None Constituents: 10% Glycerol, 20mM Tris HCl, 1mM DTT, pH 8.0
Storage	Store at +4°C short term (1-2 weeks). Aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Sequences of amino acids

MGSSHHHHHH SSGLVPRGSH MGIKVQRPRC FFDIANNQP AGRVVFELFS
 DVCPKTCENF RCLCTGEKGT GKSTQKPLHY KSCLFHRVVKDFMVQGGDFS
 EGNRGGGESI YGGFFEDES AVKHNKEFLL SMANRGKDTN GSQFFITTKP
 TPHLDGHHVV FGQVISGQEV VREIENQKTD AASKPFAEVR ILSCG

Sequence Similarities

Contains 1 PPlase cyclophilin-type domain.

GENE INFORMATION

Gene Name PPIG peptidylprolyl isomerase G (cyclophilin G) [Homo sapiens]

Official Symbol PPIG

Synonyms PPIG; peptidylprolyl isomerase G (cyclophilin G); peptidyl prolyl isomerase G (cyclophilin G); peptidyl-prolyl cis-trans isomerase G; CARS Cyp; SCAF10; SR related CTD associated factor 10; SRCyp;

Gene ID 9360

mRNA Refseq NM_004792

Protein Refseq NP_004783

MIM 606093

Uniprot ID Q13427

Chromosome Location 2q31.1

Function cyclosporin A binding; isomerase activity; peptidyl-prolyl cis-trans isomerase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA