

Active Native Wheat Germ Agglutinin Protein, Rhodamine labeled

Cat. No. Lectin-1862W **Lot. No.** (See product label)

SPECIFICATION

Product Overview	This product is the Agarose bound Wheat germ Agglutinin and has sugar specificity against N-Acetylglucosamine. The excitation maximum is at 550 nm and the emission maximum is at 575 nm.
Species	Wheat germ
Source	Wheat Germ
Description	Wheat germ agglutinin (WGA) contains a group of closely related isolectins, with an isoelectric point about pH 9. The receptor sugar for WGA is N-acetylglucosamine, with preferential binding to dimers and trimers of this sugar. WGA can bind oligosaccharides containing terminal N-acetylglucosamine or chitobiose, structures which are common to many serum and membrane glycoproteins. Bacterial cell wall peptidoglycans, chitin, cartilage glycosaminoglycans, and glycolipids can also bind WGA. Native WGA has also been reported to interact with some glycoproteins via sialic acid residues (see succinylated WGA). This lectin is used for the purification of insulin receptors and for neuronal tracing.
Form	10 mM HEPES, 0.15 M NaCl, pH 7.5, 0.08% sodium azide, 0.1 mM Ca ²⁺
Bio-activity	Inhibiting/Eluting Sugar: Chitin Hydrolysate or 500 mM N-acetylglucosamine with salt and/or acid elution generally required.
Molecular Mass	36 kDa

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications	Immunofluorescence, Glycobiology
Usage	The recommended concentration range for use is 5-20 μ g/ml.
Storage	Refrigerate in the dark. If a precipitate forms upon long-term storage at 4 centigrade, warm to 37 centigrade. and centrifuge before use.
Concentration	5 mg/ml
GENE INFORMATION	
Synonyms	Lectin; WGA

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA